


Fragrances not only scent

Pre-experimental toxicity testing for the freshwater amphipod *Hyalella azteca* bioconcentration test (HYBIT) with difficult to test chemicals

S. Groß^{1,2}, R. Schumacher², C. Mörtl², C. Schlechtriem^{1,2}

¹ RWTH Aachen University, Aachen, Germany ² Fraunhofer Institute for Molecular Biology and Applied Ecology IME, Schmallenberg, Germany



A Strong Jr, D. R. (1972). "Life history variation among populations of an amphipod (*Hyalella azteca*)."
Ecology 53(6): 1103-1111.

B Schlechtriem, C., Kampe, S., Bruckert, H. J., Bischof, I., Ebersbach, I., Kosfeld, V., ... & L'Haridon, J. (2019). Bioconcentration studies with the freshwater amphipod *Hyalella azteca*: are the results predictive of bioconcentration in fish? *Environmental Science and Pollution Research*, 26(2), 1628-1641.

C US Environmental Protection Agency. (2000). Methods for measuring the toxicity and bioaccumulation of sediment-associated contaminants with freshwater invertebrates. *Testing manual (Inland testing manual)*.

Abbreviations

- ANOVA Analysis Of VAriance
- BCF Bioconcentration Factor
- c concentration
- LC/MS Liquid Chromatography-Mass Spectrometry
- TWA Time Weighted Average
- WAF Water Accommodated Fraction

Contact:
sabine.gross@rwth-aachen.de